

UNCONTROLLABLE CRYING
and/or **LAUGHING** after
a **NEUROLOGIC CONDITION**
or a **BRAIN INJURY?**

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and
quinidine sulfate) capsules 20 mg
10 mg

What is NUEDEXTA[®] approved for?
NUEDEXTA[®] is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels. PBA is distinct and different from other types of emotional changes caused by neurologic disease or injury.

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 1

Could it be PBA
(PseudoBulbar Affect)?

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and
quinidine sulfate) capsules 20 mg
10 mg

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.
- About all medicines, herbal supplements, and vitamins you take as NUEDEXTA and certain other medicines can interact causing side effects.
- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2

Ask your doctor about
the only FDA-APPROVED
treatment for PBA

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and
quinidine sulfate) capsules 20 mg
10 mg

- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.
- If you have myasthenia gravis.

While taking NUEDEXTA, call your doctor right away:

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3

Use the
PBA CONVERSATION KIT
to start the discussion

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and
quinidine sulfate) capsules 20 mg
10 mg

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.
- If you feel dizzy, since it may increase your risk of falling.
- If you have muscle twitching, confusion, high blood pressure, fever, restlessness, sweating, or shivering, as these may be signs of a potential drug interaction called serotonin syndrome.

The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 4

BANNER 1

Frame 1	<p>UNCONTROLLABLE CRYING and/or LAUGHING after a NEUROLOGIC CONDITION or a BRAIN INJURY?</p>	<p>NUEDEXTA[®] (dextromethorphan HBr and quinidine sulfate) capsules 20 mg / 10 mg</p> <p>GET A FREE KIT</p>	<p>What is NUEDEXTA[®] approved for? NUEDEXTA[®] is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels.</p> <p>IMPORTANT SAFETY INFORMATION FULL PRODUCT INFORMATION</p>
Frame 2	<p>Could it be PBA (PseudoBulbar Affect)?</p>	<p>NUEDEXTA[®] (dextromethorphan HBr and quinidine sulfate) capsules 20 mg / 10 mg</p> <p>GET A FREE KIT</p>	<p>Important Safety Information Before you take NUEDEXTA, tell your doctor:</p> <ul style="list-style-type: none">• If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA. <p>IMPORTANT SAFETY INFORMATION FULL PRODUCT INFORMATION</p>
Frame 3	<p>Ask your doctor about the only FDA-APPROVED treatment for PBA</p>	<p>NUEDEXTA[®] (dextromethorphan HBr and quinidine sulfate) capsules 20 mg / 10 mg</p> <p>GET A FREE KIT</p>	<ul style="list-style-type: none">• If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.• If you feel dizzy, since it may increase your risk of falling. <p>IMPORTANT SAFETY INFORMATION FULL PRODUCT INFORMATION</p>
Frame 4	<p>Use the PBA CONVERSATION KIT to start the discussion</p>	<p>NUEDEXTA[®] (dextromethorphan HBr and quinidine sulfate) capsules 20 mg / 10 mg</p> <p>GET A FREE KIT</p>	<p>The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.</p> <p>You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.</p> <p>IMPORTANT SAFETY INFORMATION FULL PRODUCT INFORMATION</p>

“I wondered if I had something MORE than DEPRESSION.”

-Karen, MS Patient with PBA (PseudoBulbar Affect)



SEE REAL STORIES

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

What is NUEDEXTA® approved for?
NUEDEXTA® is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels. PBA is distinct and different from other types of emotional changes caused by neurologic disease or injury.

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 1

Karen asked her doctor about her UNCONTROLLABLE CRYING and/or LAUGHING symptoms



SEE REAL STORIES

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.
- About all medicines, herbal supplements, and vitamins you take as NUEDEXTA and certain other medicines can interact causing side effects.
- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2

Could your symptoms be PBA? Talk to your doctor about PBA and NUEDEXTA®



SEE REAL STORIES

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.
- If you have myasthenia gravis.

While taking NUEDEXTA, call your doctor right away:

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3

Frame 1



“I wondered if I had something
MORE than DEPRESSION.”

-Karen, MS Patient with PBA
(PseudoBulbar Affect)

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

What is NUEDEXTA[®] approved for?
NUEDEXTA[®] is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2



Karen asked her doctor about
her **UNCONTROLLABLE CRYING**
and/or **LAUGHING** symptoms

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3



Could your symptoms be PBA?
Talk to your doctor about
PBA and NUEDEXTA[®]

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

It's time to ask your doctor about UNCONTROLLABLE CRYING and/or LAUGHING symptoms

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

What is NUEDEXTA[®] approved for?
NUEDEXTA[®] is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels. PBA is distinct and different from other types of emotional changes caused by neurologic disease or injury.

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 1

If these symptoms followed certain NEUROLOGIC CONDITIONS or a BRAIN INJURY...

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

- NUEDEXTA is only available by prescription.

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.
- About all medicines, herbal supplements, and vitamins you take as NUEDEXTA and certain other medicines can interact causing side effects.
- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2

...ask your doctor if it could be PBA (PseudoBulbar Affect).

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.
- If you have myasthenia gravis.

While taking NUEDEXTA, call your doctor right away:

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3

Start the discussion with the PBA CONVERSATION KIT

GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg quinidine sulfate) capsules 10 mg

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.
- If you feel dizzy, since it may increase your risk of falling.
- If you have muscle twitching, confusion, high blood pressure, fever, restlessness, sweating, or shivering, as these may be signs of a potential drug interaction called serotonin syndrome.

The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 4

Frame 1

It's time to ask your doctor about **UNCONTROLLABLE CRYING** and/or **LAUGHING** symptoms

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

What is NUEDEXTA[®] approved for?
NUEDEXTA[®] is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2

If these symptoms followed certain **NEUROLOGIC CONDITIONS** or a **BRAIN INJURY...**

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3

...ask your doctor if it could be **PBA** (PseudoBulbar Affect).

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.
- If you feel dizzy, since it may increase your risk of falling.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 4

Start the discussion with the **PBA CONVERSATION KIT**

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

GET A FREE KIT

The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Do you live with
UNCONTROLLABLE CRYING
and/or **LAUGHING** symptoms?



GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

What is NUEDEXTA® approved for?
NUEDEXTA® is approved for the treatment of PseudoBulbar Affect (PBA). PBA is a medical condition that causes involuntary, sudden, and frequent episodes of crying and/or laughing in people living with certain neurologic conditions or brain injury. PBA episodes are typically exaggerated or don't match how the person feels. PBA is distinct and different from other types of emotional changes caused by neurologic disease or injury.

- NUEDEXTA is only available by prescription.


Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 1

Did they follow a
NEUROLOGIC CONDITION
or **BRAIN INJURY**?



GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

- NUEDEXTA is only available by prescription.


Important Safety Information
Before you take NUEDEXTA, tell your doctor:

- If you are taking monoamine oxidase inhibitors (MAOIs), quinidine, or quinidine-related drugs. These can interact with NUEDEXTA causing serious side effects. MAOIs cannot be taken within 14 days before or after taking NUEDEXTA.
- If you have previously had an allergic reaction to dextromethorphan, quinidine or quinidine-like drugs.
- About all medicines, herbal supplements, and vitamins you take as NUEDEXTA and certain other medicines can interact causing side effects.
- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 2

Learn how to ask
if it could be **PBA**
(PseudoBulbar Affect)



GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

- If you have had heart disease or have a family history of heart rhythm problems. NUEDEXTA may cause serious side effects, including changes in heart rhythm. If you have certain heart problems, NUEDEXTA may not be right for you. Your doctor may test your heart rhythm (heartbeats) before you start NUEDEXTA.
- If you have myasthenia gravis.


While taking NUEDEXTA, call your doctor right away:

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 3

The **PBA Conversation**
Kit can help you
talk to your doctor



GET A FREE KIT

NUEDEXTA
(dextromethorphan HBr and 20 mg
quinidine sulfate) capsules 10 mg

- If you feel faint or lose consciousness.
- If you experience lightheadedness, chills, fever, nausea, or vomiting as these may be signs of an allergic reaction to NUEDEXTA. Hepatitis has been seen in patients taking quinidine, an ingredient in NUEDEXTA.
- If you have unexplained bleeding or bruising. Quinidine, an ingredient in NUEDEXTA, can cause a reduction in the number of platelets in your blood which can be severe and, if left untreated, can be fatal.
- If you feel dizzy, since it may increase your risk of falling.
- If you have muscle twitching, confusion, high blood pressure, fever, restlessness, sweating, or shivering, as these may be signs of a potential drug interaction called serotonin syndrome.

The most common side effects of NUEDEXTA are diarrhea, dizziness, cough, vomiting, weakness, and swelling of feet and ankles. This is not a complete list of side effects. Tell your doctor about any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 800-FDA-1088.

[IMPORTANT SAFETY INFORMATION](#) | [FULL PRODUCT INFORMATION](#)

Frame 4

BANNER 5




Save on
NUEDEXTA[®]
PRESCRIPTIONS
and REFILLS

[DOWNLOAD A CARD](#)

NUEDEXTA[®]
(dextromethorphan HBr and $\frac{20\text{ mg}}$
quinidine sulfate) capsules $\frac{10\text{ mg}}$

©2019 Avanir Pharmaceuticals, Inc. All rights reserved.
AVANIR and NUEDEXTA are trademarks or registered trademarks
of Avanir Pharmaceuticals, Inc. in the United States and other countries. MLR-NUE-US-XXXX-XXXX

Frame 1



Pay
AS LITTLE AS \$0
for a **90-DAY**
prescription*


*RESTRICTIONS APPLY

[DOWNLOAD A CARD](#)

NUEDEXTA[®]
(dextromethorphan HBr and $\frac{20\text{ mg}}$
quinidine sulfate) capsules $\frac{10\text{ mg}}$

©2019 Avanir Pharmaceuticals, Inc. All rights reserved.
AVANIR and NUEDEXTA are trademarks or registered trademarks
of Avanir Pharmaceuticals, Inc. in the United States and other countries. MLR-NUE-US-XXXX-XXXX

Frame 2



Or pay
AS LITTLE AS \$30
for a **30-DAY**
prescription*


*RESTRICTIONS APPLY

[DOWNLOAD A CARD](#)

NUEDEXTA[®]
(dextromethorphan HBr and $\frac{20\text{ mg}}$
quinidine sulfate) capsules $\frac{10\text{ mg}}$

©2019 Avanir Pharmaceuticals, Inc. All rights reserved.
AVANIR and NUEDEXTA are trademarks or registered trademarks
of Avanir Pharmaceuticals, Inc. in the United States and other countries. MLR-NUE-US-XXXX-XXXX

Frame 3



Get your
Co-Pay
SAVINGS CARD
now

*RESTRICTIONS APPLY

[DOWNLOAD A CARD](#)

NUEDEXTA[®]
(dextromethorphan HBr and $\frac{20\text{ mg}}$
quinidine sulfate) capsules $\frac{10\text{ mg}}$

©2019 Avanir Pharmaceuticals, Inc. All rights reserved.
AVANIR and NUEDEXTA are trademarks or registered trademarks
of Avanir Pharmaceuticals, Inc. in the United States and other countries. MLR-NUE-US-XXXX-XXXX

Frame 4